

EDITORIAL REVIEW

Antigen shedding and metastasis of tumour cells

S. K. A. LAW MRC Immunochemistry Unit, Department of Biochemistry, University of Oxford, Oxford, England

(Accepted for publication 19 April 1991)

Release of membrane surface antigens appears to be a common feature of malignant cells (Black, 1980; Baker *et al.*, 1982). The antigens have been detected in culture media of active growing cell lines *in vitro* (Komada *et al.*, 1987; Chiba *et al.*, 1989) and in the body fluids of humans and animals with various malignant tumours *in vivo* (Komada *et al.*, 1990; Vinuela *et al.*, 1991). The antigens detected were not confined to any particular molecule: they include the MHC antigens (Tsujisaki *et al.*, 1987), other differentiation antigens (Kay *et al.*, 1988; Komada *et al.*, 1990), tumour-specific/associated antigens (Fauget & Agee, 1989; Chiba *et al.*, 1989), and, in this issue of *Clinical and Experimental Immunology*, the intercellular adhesion molecule-1 (ICAM-1) (Tsujisaki *et al.*, 1991). In the case of cultured cells, the amount of antigen shed appears to correlate with cell growth and not cell division or cell death (Chiba *et al.*, 1989) whereas in patients, the level of antigens in bodily fluids seems to correlate with the clinical stage of the tumour (Fauget & Agee, 1989; Komada *et al.*, 1990).

When a cell is engaged in active growth, the general increase in cellular content, including molecules that constitute the membrane, may be taken as proportional to the increase in cell volume. The increase in membrane components is therefore in disproportionate excess compared with the increase in cell surface area, which varies, with approximation, as the 2/3rd power of the cell volume on the assumption that some simple balance of cell volume and cell surface area is to be maintained. The excess membrane must be disposed of, and one of the possible means is the shedding in the form of vesicles. Indeed, extensive shedding had been observed in different tumour cell lines by electron microscopy. Vesicles were either recovered in the pellets upon ultracentrifugation of tissue culture supernates and ascites fluid (Dvorak *et al.*, 1981), or observed as out-growths from the parent cell body (Chiba *et al.*, 1989). In addition, the shedding of vesicles from the rat fibrosarcoma (Chiba *et al.*, 1989) was observed when cultured in 10% fetal calf serum (FCS) but not when under less encouraging growth conditions of 1% FCS. Other mechanisms of antigen release have also been suggested, including their proteolysis and the secretion of variant forms (Kay *et al.*, 1988).

ICAM-1 was first described by Rothlein *et al.* (1986) as an adhesion molecule that participates in leucocyte function-associated antigen-1 (LFA-1) dependent homotypic adherence

of phorbol-ester-activated leucocytes and was subsequently established to be the counter-receptor for LFA-1 (Martin & Springer, 1987). The LFA-1/ICAM-1 interaction is important in a number of leucocyte adhesion activities (Makgoba *et al.*, 1988a) including the conjugate formation between cytolytic T cell and their targets (Makgoba *et al.*, 1988b) and natural killer (NK) mediated cytotoxicity (Kohl *et al.*, 1984; Timonen, Patarroyo & Gahmberg, 1988). It is generally recognized that NK cells are involved in the surveillance and regulation of tumour metastasis (Herberman & Ortaldo, 1981; Storkus & Dawson, 1991): the metastatic potential of tumour cells, in their blood-borne phase, shows a negative correlation with their sensitivity to NK cytotoxicity (Kawano *et al.*, 1986; Greenberg *et al.*, 1987) and the depletion of NK cells in mice results in enhanced susceptibility of the animals to malignancy (Hanna & Burton, 1981; Kawase *et al.*, 1982; Wiltout *et al.*, 1985; Seaman *et al.*, 1987). Thus, the expression and shedding of ICAM-1 by tumour cells, as reported by Tsujisaki *et al.* (1991) in this issue, may have profound implications for the metastasis of tumour cells and for its regulation by cells of the immune system in terms of the tumour cells escaping surveillance by the shedding of certain membrane antigens.

The expression of ICAM-1 on tumour cells is stimulated by interferon-gamma (IFN- γ), and perhaps other cytokines, probably as a result of some form of tumour-induced T cell activation. The immediate consequence would be the enhancement of sensitivity of the tumour cells to LFA-1/ICAM-1-dependent adhesion and cytotoxicity. The increase of expression of ICAM-1 on fibroblasts surrounding the tumour nests may promote the adhesion NK cells to fibroblasts. Since this interaction is probably insufficient to induce NK cytotoxicity of fibroblasts, it is transient and may serve to direct the migration of NK cells towards the tumour target. If the tumour cells, or a sufficient number of them, survive this phase of surveillance, active growth may lead to the shedding of ICAM-1-containing membrane vesicles which, by interacting with LFA-1 on NK and other effector cells, would revert the balance to favour the survival and further growth, and eventually the metastasis, of the tumour cells. This view is consistent with the observation that IFN treatment of tumour cells can provide protection against NK cytotoxicity and enhance their metastatic capacity (Welsh *et al.*, 1981; McMillan *et al.*, 1987).

Other antigens and molecules in the vesicles shed from tumour cells may also play a role in their survival. Disproportionate shedding of tumour antigen may deplete the parent cell of tumour antigen (Chiba *et al.*, 1989), thus making them invisible

Correspondence: S. K. A. Law, MRC Immunochemistry Unit, Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU, UK.

to the host immune surveillance system. Alternatively, tumour-specific antigens on shed vesicles can absorb circulating antibodies thus neutralizing their anti-tumour effects (Vinuela *et al.*, 1991). Promoters of coagulation have been detected in shed vesicles; it is argued that they may induce clotting in the proximity of the tumour, thus forming a barricade to isolate the tumour from cells of the host immune system (Dvorak *et al.*, 1981).

The measurement of the level of shed antigens in body fluids may provide a convenient means to monitor the clinical state of cancer patients. Such a correlation had been documented for CD9 in the cerebrospinal fluid of patients with acute lymphoblastic leukaemia (Komada *et al.*, 1990), and for IL-2 receptors as well as tumour-associated antigens in the serum of patients with B-chronic lymphocytic leukaemia (Kay *et al.*, 1988; Fauget & Agee, 1989). It now appears that the presence of circulating ICAM-1 may provide a general indication of malignant disease incidents.

REFERENCES

- BAKER, M.A., RONCARI, D.A.K., TAUB, R.N., MOHANAKUMAR, T., FALK, J.A. & GRANT, S. (1982) Characterization of compounds shed from the surface of human leukemic myeloblasts in vitro. *Blood*, **60**, 412.
- BLACK, P.H. (1980) Shedding from the cell surface of normal and cancer cells. *Adv. Cancer. Res.* **32**, 75.
- CHIBA, I., JIN, R., HAMADA, J., HOSOKAWA, M., TAKEICHI, N. & KOBAYASHI, H. (1989) Growth-associated shedding of a tumor antigen (CE7) detected by a monoclonal antibody. *Cancer Res.* **49**, 3972.
- DVORAK, H.F., QUAY, S.C., ORENSTEIN, N.S., DVORAK, A.M., HAHN, P., BITZER, A.M. & CARVALHO, A.C. (1981) Tumor shedding and coagulation. *Science*, **212**, 923.
- FAUGET, G.B. & AGE, J.F. (1989) Modulation, shedding, and serum titers of the chronic lymphatic leukemia-associated antigen: characterization and clinical correlations. *Blood*, **74**, 2493.
- GREENBERG, A.H., EGAN, S.E., JAROLIM, L., GINGRAS, M.C. & WRIGHT, J.A. (1987) Natural killer cell regulation of implantation and early lung growth of H-ras-transformed 10T1/2 fibroblasts in mice. *Cancer Res.* **47**, 4801.
- HANNA, N. & BURTON, R.C. (1981) Definitive evidence that natural killer (NK) cells inhibit experimental tumor metastasis in vivo. *J. Immunol.* **127**, 1754.
- HERBERMAN, R.B. & ORTALDO, J.R. (1981) Natural killer cells: their role in defense against disease. *Science*, **214**, 24.
- KAWANO, Y.I., TANIGUCHI, K., TOSHITANI, A. & NOMOTO, K. (1986) Synergistic defense system by cooperative natural effectors against metastasis of B16 melanoma cells in H-2-associated control: different behavior of H-2⁺ and H-2⁻ cells in metastatic processes. *J. Immunol.* **136**, 4729.
- KAWASE, I., URDAL, D.L., BROOKS, C.G. & HENNEY, C.S. (1982) Selective depletion of NK cell activity in vivo and its effects on the growth of NK-sensitive and NK-resistant tumor cell variants. *Int. J. Cancer*, **29**, 567.
- KAY, N.E., BURTON, J., WAGNER, D. & NELSON, D.L. (1988) The malignant B cells from B-chronic lymphocytic leukemia patients release TAC-soluble interleukin-2 receptors. *Blood*, **72**, 447.
- KOHL, S., SPRINGER, T.A., SCHMALSTIEG, F.C., LOO, L.S. & ANDERSON, D.C. (1984) Defective natural killer cytotoxicity and polymorphonuclear leukocyte antibody-dependent cellular cytotoxicity in patients with LFA-1/QKM1 deficiency. *J. Immunol.* **133**, 2972.
- KOMADA, Y., OCHIAI, H., SHIMIZU, K., AZUMA, E., KAMIYA, H. & SAKURAI, M. (1987) Shedding of leukemia-associated P24 antigen by lymphoblastoid cell lines. *Jpn. J. clin. Oncol.* **17**, 333.
- KOMADA, Y., OCHIAI, H., SHIMIZU, K., AZUMA, E., KAMIYA, H. & SAKURAI, M. (1990) Shedding of CD9 antigen into cerebrospinal fluid by acute lymphoblastic leukemia cells. *Blood*, **76**, 112.
- McMILLAN, T.J., RAO, J., EVERETT, C.A. & HART, I.R. (1987) Interferon-induced alterations in metastatic capacity, class-I antigen expression, and natural killer cell sensitivity of melanoma cells. *Int. J. Cancer*, **40**, 659.
- MAKGOBA, M.W., SANDERS, M.E., GINTHER-LUCE, G.E., DUSTIN, M.L., SPRINGER, T.A., CLARK, E.A., MANNONI, P. & SHAW, S. (1988a) ICAM-1 a ligand for LFA-1-dependent adhesion of B, T and myeloid cells. *Nature*, **331**, 86.
- MAKGOBA, M.W., SANDERS, M.E., GINTHER-LUCE, G.E., GUGEL, E.A., DUSTIN, M.L., SPRINGER, T.A. & SHAW, S. (1988b) Functional evidence that intercellular adhesion molecule-1 (ICAM-1) is a ligand for LFA-1-dependent adhesion in T cell-mediated cytotoxicity. *Eur. J. Immunol.* **18**, 637.
- MARLIN, S.D. & SPRINGER, T.A. (1987) Purified intercellular adhesion molecule-1 (ICAM-1) is a ligand for lymphocyte function-associated antigen 1 (LFA-1) *Cell*, **51**, 813.
- ROTHLEIN, R., DUSTIN, M.L., MARLIN, S.D. & SPRINGER, T.A. (1986) A human intercellular adhesion molecule (ICAM-1) distinct from LFA-1. *J. Immunol.* **137**, 1270.
- SEAMAN, W.E., SLEISINGER, M., ERIKSSON, E. & KOO, G.C. (1987) Depletion of natural killer cells in mice by monoclonal antibody to NK-1.1. Reduction in host defense against malignancy without loss of cellular or humoral immunity. *J. Immunol.* **138**, 4539.
- STORKUS, W.J. & DAWSON, J.R. (1991) Target structures involved in natural killing (NK): characteristics, distribution, and candidate molecules. *CRC crit. Rev. Immunol.* **10**, 393.
- TIMONEN, T., PATARROYO, M. & GAHMBERG, C.G. (1988) CD11a-c/CD18 and GP84 (LB-2) adhesion molecules on human large granular lymphocytes and their participation in natural killing. *J. Immunol.* **141**, 1041.
- TSUJISAKI, M., IGARASHI, M., SAKAGUCHI, K., EISINGER, M., HERLYN, M. & FERRONE, S. (1987) Immunochemical and functional analysis of HLA class II antigens induced by recombinant immune interferon on normal epidermal melanocytes. *J. Immunol.* **138**, 1310.
- TSUJISAKI, M., IMAI, K., HIRATA, H., HANZAWA, Y., MASUYA, J., NAKANO, T., SUGIYAMA, T., MATSUI, M., HINODA, Y. & YACHI, A. (1991) Detection of circulating intercellular adhesion molecule-1 antigen in malignant diseases. *Clin. exp. Immunol.* **85**, 3.
- VINUELA, J.E., RODRIGUEZ, R., GIL, J., COLL, J., DE LA CONCHA, E.G. & SUBIZA, J.L. (1991) Antigen shedding vs development of natural suppressor cells as mechanism of tumor escape in mice bearing Ehrlich tumor. *Int. J. Cancer*, **47**, 86.
- WELSH, R.M., KARRE, K., HANSSON, M., KUNKEL, L.A. & KIESSLING, R.W. (1981) Interferon-mediated protection of normal and tumor target cells against lysis by mouse natural killer cells. *J. Immunol.* **126**, 219.
- WILTROUT, R.H., HERBERMAN, R.B., ZHANG, S.R., CHIRIGOS, M.A., ORTALDO, J.R., GREEN, K.M. & TALMADGE, J.E. (1985) Role of organ-associated NK cells in decreased formation of experimental metastases in lung and liver. *J. Immunol.* **134**, 4267.